



THE PRACTITIONER LE PRATICIEN

The occasional management of epistaxis

Todd K. Young, BScN,
MD, CCFP

Green Bay Health Centre,
Springdale, NL
Memorial University of
Newfoundland, St. John's,
NL
Northern Ontario School of
Medicine, Thunder Bay, Ont.

Ryan Hall, BScN
Northern Ontario School of
Medicine, Thunder Bay, Ont.

Correspondence to:
Dr. Todd Young, Box 1705,
Springdale NL A0J 1T0;
tkyoung@explornet.com

This article has been peer
reviewed.

INTRODUCTION

Epistaxis, or nasal bleeding, is the most common otolaryngologic emergency, reported to occur in up to 60% of the general population.^{1,2} Although the presentation of epistaxis can be quite distressing to the patient and physician, only 10% of adult patients with epistaxis require definitive medical attention.³ In rare cases, however, massive nasal bleeding can be fatal.⁴

ANATOMY

Knowledge of nasal anatomy is essential in the detection of the bleed location and subsequent hemostasis. Although nasal circulation is complex (Fig. 1), epistaxis is usually described as either anterior or posterior. This distinction provides a useful basis for management.²

ANTERIOR BLEEDS

Anterior bleeds are by far the most common.⁵ Up to 80% occur within the




Fig. 1. Cross section of normal nasal circulation. © 2004 Christy Krames with permission.

clinically obvious vascular watershed area of the nasal septum known as the Kiesselbach plexus.⁶ Anterior nose-bleeds often result from mucosal trauma or irritation such as the following:⁵

- nose picking (most common)
- facial trauma secondary to motor vehicle collision or other blunt facial impact
- mucosal hyperemia secondary to allergic or viral rhinitis
- presence of a foreign body (if bleed is accompanied by purulent discharge)
- chronic excoriation secondary to chronic intranasal drug use

POSTERIOR BLEEDS

Posterior epistaxis can result in significant hemorrhage⁵ and generally arises from the posterior nasal cavity via branches of the sphenopalatine arteries.⁷ Posterior epistaxis occasionally may be asymptomatic or may present insidiously as nausea, hematemesis, anemia, hemoptysis or melena.²

RISK FACTORS

Risk factors that increase the incidence of anterior and posterior epistaxis include the following:

- anticoagulation⁸
- hereditary hemorrhagic telangiectasia or Osler-Weber-Rendu disease (most common presenting symptom of this disease)⁹
- blood dyscrasias, particularly platelet disorders, von Willebrand disease and hemophilia⁵
- aneurysms of the head and neck vasculature secondary to prior regional surgery⁶

- nasal neoplasms
- chronic alcohol abuse¹⁰
- intranasal steroid use¹¹

The associations regarding use of acetylsalicylic acid and hypertension as risk factors for epistaxis are uncertain.^{2,12}

EQUIPMENT

An epistaxis tray can be created using common supplies and a few specialized instruments:²

- nasal decongestant spray
- local anesthetic
- silver nitrate cautery sticks
- suction cautery (if available)
- bayonet forceps
- nasal speculum
- lubricating jelly
- bacitracin, mupirocin or other antistaphylococcal ointment¹³
- Frazier suction tip or suction tube connector (which works nicely as a suction tip in this situation)¹⁴
- posterior double balloon system or 14-F Foley catheter and syringe for balloon inflation
- packing materials including Merocel packs (Medtronic Inc.), Gelfoam (Pfizer, Inc.) and Xeroform (Kendall Healthcare; nonadherent gauze impregnated with petroleum jelly and 3% tribromophenate)
- emesis basin

MANAGEMENT

Step 1: evaluation and stabilization

- Initial management should focus on the cardiovascular stability of the patient. Massive epistaxis may necessitate airway intervention and fluid resuscitation before hemostasis is attempted. Normal appearance, vital signs and respiratory function are evidence that you can safely attend to the epistaxis.⁵
- Consider intravenous access. Difficult cases may require anxiolytic and/or analgesic medications.²
- Epistaxis can be a very messy situation. Always remember universal precautions and the “3 Gs”: gloves, gown and goggles.¹⁵
- Take a quick but comprehensive history (Box 1).¹⁴
- The patient should be in a well-lit room, seated upright, looking directly ahead and attempting the sniffing position. This allows for optimal view of the nasopharynx.⁵

- Clots may be cleared via suction or by asking the patient to *gently* blow his or her nose.⁵
- Attempt to visualize the bleed. The Kiesselbach plexus should be examined first since most bleeds originate here.⁵

Step 2: direct pressure

- Ask the patient to apply direct pressure by pinching the lower part of the nose (not the bridge) for 10 to 15 minutes.¹⁴
- The patient should be mouth breathing and leaning forward.
- Encourage the patient to spit out posterior pharyngeal blood into a basin. This will reduce the risk of vomiting from swallowed blood and of aspiration.⁵
- An ice pack over the dorsum of the nose may assist with hemostasis by constricting the blood vessels and thereby stopping the bleeding. Applying an ice pack to the back of the neck activates the mammalian dive reflex, thereby causing peripheral vasoconstriction.¹³
- Consider spraying the nasal cavity with decongestant spray, reapplying nasal pressure and waiting a few minutes.^{5,14}
- An alternate method is to insert cotton pledgets, soaked with an anesthetic–vasoconstrictor solution, into the nasal cavity to anesthetize and shrink nasal mucosa. Soak pledgets in 4% topical cocaine solution or a solution of 4% lidocaine and topical epinephrine (1:10 000) and place them into the nasal cavity. Leave them in place for 10–15 minutes.¹⁶
- Most anterior nosebleeds stop with the above procedures. If this does not occur, it is time to proceed to cauterizing the bleeding points.¹³

Step 3: cautery

- Electrical or chemical cautery can be used if an anterior source is identified. Chemical cautery is

Box 1. Factors to consider when taking the history of the patient with epistaxis

- Which side is bleeding?
- Amount of blood loss
- Is it recurrent?
- Is it in the pharynx?
- Recent trauma?
- Symptoms of hypovolemia?
- Medical history and current medication (e.g., acetylsalicylic acid, warfarin)

usually accomplished with silver nitrate sticks; this procedure will be described below.⁵

- Reinspect the nasal cavity and suction out any blood or clots to attempt to revisualize the bleeding sites.¹⁴
- Adequate illumination is important. If a headlamp is not available, an otoscope can be used.¹⁴
- Once the bleeding site has been identified, the silver nitrate applicator tip is applied to a small area surrounding the bleeding site starting proximally and moving centrally in a radial fashion. Use the nasal speculum, and brace your cauterizing hand. Ask your patient to remain still and explain why.^{5,14}
- Silver nitrate cauterizes everything it touches. Do not touch the facial skin, nasal alae or other nasal mucosa. In addition, silver nitrate may not work if bleeding is active.¹⁴
- Apply cautery for no more than 10 to 20 seconds at each site. Overzealous cautery can lead to ulceration and perforation.⁵
- To avoid the increased risk of tissue necrosis, never cauterize both sides of the septum during the same session.⁵
- If there is no more active bleeding, the patient may be discharged with printed instructions, provided there are no other medical problems.¹⁴

Step 4A: anterior packing

If bleeding continues, consider nasal packing.

Merocel packing

- Coat tampon with lubricating jelly, antibiotic ointment or a water-based cream (possibly

decreases the risk of toxic shock syndrome).⁵

- Insert the device directly along the floor of the nasal cavity (Fig. 2). The device can be re-expanded with 10 to 20 mL of saline. A 22-gauge angiocatheter on a saline syringe can be used to apply a small amount of saline posteriorly first to speed diffusion.^{5,14}
- Ensure that the tampon is inserted completely.
- Tape the strings to each cheek and apply a nasal bolster.

Xeroform or ribbon gauze packing

- Pick up the gauze with the forceps about 10–15 cm from the tip.
- Pack the nasal cavity starting from the floor of the nose upward.
- Continue to move upward by layering the gauze until you reach the cavity roof. Pack the cavity tightly.
- Both ends of the ribbon gauze should protrude from the nostril. You don't want one end dangling down the nasopharynx and causing the patient to gag, or worse, eliciting laryngospasm.
- Secure the gauze. Apply a nasal bolster to control dripping and secretions.¹⁴

Continued bleeding

If bleeding persists despite the initial packing, the contralateral naris may be packed next, providing a counterforce to promote tamponade. If this manoeuvre still does not produce hemostasis, the odds of a posterior source increase greatly, since nasal packing in anterior bleeding has about a 90%–95% success rate.^{17,18}




Fig. 2. Initial insertion angle of the Merocel pack (A). Advance the Merocel pack horizontally (B). Make certain that the pack is advanced fully into the nose (C).

Step 4B: posterior packing

- Continued hemorrhage despite an anterior pack may be due to a posterior bleed.
- Many postnasal packs are available commercially; however, the most commonly used is a Foley catheter.
- Before proceeding with the posterior packing, remove the anterior pack and re-examine the nasal cavity.
- Suction as required.⁵

Foley catheter

- Cut the tip of a 14-F Foley to minimize irritation of the posterior structures.
- Using a pair of forceps, advance the catheter along the floor of the nose until it is visible in the mouth. Partially fill the balloon with sterile saline (5–7 mL), and retract the catheter, and the cotton pack, until it is well opposed to the posterior nasopharynx.
- Fill the balloon completely (another 5 mL).
- Pain or distention of the soft palate suggests overfilling.
- At this point, consider repacking the anterior nose. Many practitioners will still pack the anterior nose at this point because without perfect opposition, some blood will pool anteriorly and exit the naris. Additionally, some epistaxis episodes involve both anterior and posterior sources, especially in the setting of a coagulopathy or hereditary hemorrhagic telangiectasia.
- Clamp the catheter in place with an umbilical clamp or a small c-clamp, as from a nasogastric tube.
- Ensure that the catheter is not pressing on the nose as alar necrosis may occur. This can be prevented by applying gauze around the nostril.⁵

Specialized balloon systems

- There are many balloon systems available. Most are effective for managing posterior bleeding.
- If you are using a double-balloon device, pass it into the affected nostril until it reaches the nasopharynx.
- Inflate the posterior balloon with 7 to 10 mL of saline, and carefully withdraw the catheter extending out of the nostril so that the balloon seats in the posterior nasal cavity to tamponade the bleeding source.
- Next, inflate the anterior balloon with roughly

15 to 30 mL of saline in the anterior nasal cavity to prevent retrograde travel of the posterior balloon and subsequent airway obstruction.

- An umbilical clamp can be placed across the stalk of the balloon adjacent to the nostril to further prevent dislodgement; the clamp should be padded to prevent pressure necrosis of the nasal skin.
- Balloon packs generally are left in place for 48–72 hours. As with anterior packing, tissue necrosis can occur if a posterior pack is inserted improperly or balloons are overinflated.²

Persistent bleeding despite anterior and posterior packing

- Patients with anterior or posterior bleeding that continues despite packing and balloon procedures should be referred to an otolaryngologist.^{2,5}
- Endoscopy can be used to locate the site of bleeding and perform direct cauterization.

Step 5: Consider antibiotics for prophylaxis of toxic shock syndrome

- The incidence of toxic shock syndrome with nasal packing is estimated at about 16 per 100 000 post-operative packings, but the incidence in primary nasal packing is not established.⁵
- It is unclear whether oral antibiotics are required for the prophylaxis of toxic shock syndrome. However, if antibiotics are prescribed, an antibiotic with staphylococcal coverage should be selected, such as amoxicillin clavulanate or a second-generation cephalosporin.^{13,19}

INVESTIGATIONS

It is recommended that coagulation screening should be carried out only if firm indications are found in the history and physical examination of patients presenting to the emergency department with epistaxis. Coagulation screening of all patients who present with epistaxis is of little value and leads to longer stays in the emergency department. Coagulation studies are justified in patients receiving anticoagulant treatment and in those with known coagulopathy or chronic liver disease.²⁰

SUMMARY

Epistaxis is a common clinical condition. It is important for clinicians to develop an approach to diagnosis and management, and it is important to distinguish between anterior and posterior bleeds. The

stepwise approach we have highlighted identifies the necessary equipment vital for timely and effective management. Clinicians are advised to prevent complications by using appropriate equipment, prescribing prophylactic antibiotics when needed, and, when bleeding continues despite packing and balloon procedures, referring patients to an otolaryngologist.

REFERENCES

1. Douglas R, Wormald PJ. Update on epistaxis. *Curr Opin Otolaryngol Head Neck Surg* 2007;15:180-3.
2. Kucik CJ, Clenney T. Management of epistaxis. *Am Fam Physician* 2005;71:305-11.
3. Petruson B, Rudin R. The frequency of epistaxis in a male population sample. *Rhinology* 1975;13:129-33.
4. Tan LK, Calhoun KH. Epistaxis. *Med Clin North Am* 1999;83:43-56.
5. Alter H. Approach to the adult with epistaxis. Available: www.utdol.com/patients/content/topic.do?topicKey=~2056/corp_da3 (accessed 2008 Dec. 29).
6. Alvi A, Joyner-Triplett N. Acute epistaxis. How to spot the source and stop the flow. *Postgrad Med* 1996;83-90;94-96.
7. Koh E, Frazzini VI, Kagetsu NJ. Epistaxis: vascular anatomy, origins, and endovascular treatment. *AJR Am J Roentgenol* 2000;174:845-51.
8. Nitu IC, Perry DJ, Lee CA. Clinical experience with the use of clotting factor concentrates in oral anticoagulation reversal. *Clin Lab Haematol* 1998;20:363-7.
9. Shah RK, Dhingra JK, Shapshay SM. Hereditary hemorrhagic telangiectasia: a review of 76 cases. *Laryngoscope* 2002;112:767-73.
10. McGarry GW, Gatehouse S, Hinnie J. Relation between alcohol and nose bleeds. *BMJ* 1994;309:640.
11. Keith P, Nieminen J, Hollingworth K, et al. Efficacy and tolerability of fluticasone propionate nasal drops 400 microgram once daily compared with placebo for the treatment of bilateral polyposis in adults. *Clin Exp Allergy* 2000;30:1460-8.
12. Kotecha B, Fowler S, Harkness P, et al. Management of epistaxis: a national survey. *Ann R Coll Surg Engl* 1996;78:444-6.
13. El-Guindy A. Endoscopic transseptal sphenopalatine artery ligation for intractable posterior epistaxis. *Ann Otol Rhinol Laryngol* 1998;107:1033-7.
14. Toner JG, Walby AP. Comparison of electro and chemical cautery in the treatment of anterior epistaxis. *J Laryngol Otol* 1990;104:617-8.
15. Leong SCL, Roe RJ, Karkanevatous A. No frills management of epistaxis. *Emerg Med J* 2005;22:470-2.
16. Bamimore O, Silverberg MA. Epistaxis. Available: www.emedicine.com/EMERG/topic806.htm (accessed 2008 Dec. 29).
17. Corbridge RJ, Djazaeri B, Hellier WP, et al. A prospective randomized controlled trial comparing the use of merocel nasal tampons and BIPP in the control of acute epistaxis. *Clin Otolaryngol* 1995;20:305-7.
18. McGlashan JA, Walsh R, Dauod A, et al. A comparative study of calcium sodium alginate (Kaltostat) and bismuth tribromophenate (Xeroform) packing in the management of epistaxis. *J Laryngol Otol* 1992;106:1067-71.
19. Derkay CS, Hirsch BE, Johnson JT, et al. Posterior nasal packing. Are intravenous antibiotics really necessary? *Arch Otolaryngol Head Neck Surg* 1989;115:439-41.
20. Awan MS, Iqbal M, Imam SZ. Epistaxis: When are coagulation studies justified? *Emerg Med J* 2008;25:156-7.

RURALMED: THE SRPC LISTSERV

MEDRURALE : LA LISTE DE DIFFUSION DE LA SMRC

RURALMED

Subscription to RuralMed is by request. Send an email message to: admin@srpc.ca.

Include your full name and email address. If you include a short biography it will be posted to the list as your introduction. You can also access both the RuralMed archives and a RuralMed subscription form through the SRPC home page at: www.srpc.ca.

MEDRURALE

Pour vous abonner au serveur de liste francophone, MedRurale, veuillez envoyer un courriel à l'adresse suivante : lamarche@comnet.ca.

Donner votre nom au complet et votre adresse de courriel. Si vous ajoutez aussi une courte biographie, elle pourra être affichée sur la liste en guise de présentation. Vous pouvez aussi accéder aux archives de MedRurale et à un formulaire d'inscription au serveur de liste anglophone sur la page d'accueil du site de la SCMR, www.srpc.ca.